

Rossby waves: Synergy in action

Graham D. Quartly, Paolo Cipollini, David Cromwell & Peter G. Challenor

Laboratory for Satellite Oceanography

Southampton Oceanography Centre

Empress Dock, Southampton,

Hants, SO14 3ZH, UK

Summary

Rossby waves are an important phenomenon linking processes in the west of ocean basins with forcing that occurred in the east. We show evidence for such features in satellite-derived datasets of sea surface height, temperature and ocean colour, using a section of the south Indian Ocean as an example. We discuss the possible mechanisms for an effect on chlorophyll, and investigate this by comparing the ocean colour signal with the other datasets. In this region, the primary mechanism for a Rossby wave signal in ocean colour appears to be advection, however this cannot fully explain the observations in the westernmost basin.

Keywords

Rossby waves, Sea surface height, Sea surface temperature, Chlorophyll concentration, Indian Ocean, Synergy

1. Introduction

Rossby waves are an important process by which signals are transmitted from one side of an ocean basin to another. They correspond to a solution of the equations of motion of a fluid on a rotating body in response to some initial perturbation, such as marked wind bursts. Oceanic Rossby waves have a large wavelength (of order 1000 km) and slow propagation (of order 1-10 km day⁻¹). They are not just a surface phenomenon, but may affect the full depth of the water column, and have the intriguing property that for long wavelengths the group velocity is predominantly westward. This means that they can only convey information (on changes in forcing) from east to west. Further details on the transmission and vertical structure of Rossby waves can be found in Gill (1982). Although the concept of Rossby waves and their observation in the atmosphere have been known for more than half a century, there were limited observations of oceanic Rossby waves until the advent of satellite sensors able to provide accurate basin-wide observations every few days. In this short paper we illustrate the observations of Rossby waves in sea surface height, temperature and ocean colour datasets, and discuss the challenges in interpretation that have arisen.

2. Satellite observations and filtering

Tokmakian and Challenor (1993) first showed that Rossby waves could be detected from space. Subsequently, Chelton and Schlax (1996) have demonstrated that Rossby waves are present in all ocean basins at many different latitudes. Hill et al. (2000) were the first to show the near-ubiquity of these signals in SST and even more recently Cipollini et al. (2001) and Uz et al. (2001) noted the chlorophyll signature of Rossby

waves in all the main ocean basins. To illustrate the signature of Rossby waves within various datasets we show longitude-time sections (Fig. 1) of sea surface height (SSH), sea surface temperature (SST) and chlorophyll concentration (CC) at 34°S in the Indian Ocean, as this is a location where the signals for all three are particularly clear.

For the temperature dataset we use measurements from the Along-Track Scanning Radiometer (ATSR-2), which has a stated accuracy for point measurements of 0.3K (Mutlow et al., 1994). Although a Hovmöller plot of SST (Fig. 1a) is dominated by the seasonal cycle of warming and cooling, hints of westward-propagating features can be discerned. To enhance these we apply a high-pass filter in the spatial domain. This is simply a matter of differencing the value at a location with respect to a 550-km running mean. This enhances the shorter scale features at the expense of the basin-wide consistent changes (see Fig. 1b). The typical amplitude of these resolved features is 0.5K, with positive and negative anomalies being equally pronounced.

SeaWiFS provides information on ocean colour, from which chlorophyll concentration can be calculated. The stated accuracy for 1km² pixels is of order 30% (Hooker and McClain, 2000), but, provided some of the errors are not systematic, the overall uncertainty of the monthly averages used here will be smaller. As the distribution of CC values is approximately log-normal (Campbell, 1995), it is sensible to perform our analysis on the logarithms of the values. This means that our filtering returns the *ratio* of CC at a location to the value of the running mean. Again westward features can be discerned, with their peak amplitude being in December-January, and the peak occurring slightly earlier in the west than the east. The peak in the CC signal of Rossby waves occurs at the *end* of the spring bloom, which occurs earlier in the west than the east. Although the amplitude of the chlorophyll signature of Rossby waves is at times only 10% (e.g. July-August in the east), there is still a coherent pattern of propagation.

Finally in Fig. 1d we provide the SSH signature of Rossby waves as detected by the TOPEX/Poseidon altimeter. The altimetric dataset usually contains the clearest indicator of Rossby waves, partly because SSH is an integrated quantity over the entire water column, rather than a surface manifestation that can be masked by local conditions. The surface amplitude here of Rossby waves is of order 5cm, which is significantly above the 2cm measurement error of the instrument (Cheney et al., 1994).

3. Correlated observations

Westward-propagating Rossby waves involve the meridional displacements of water. Gill (1982) explains how this relates to changes in currents and thus a SSH signal. In the "waveguide" at 34°N in the North Atlantic, Cipollini et al. (1997) noted that the SSH and SST signatures of Rossby waves were in phase, which was explained in terms a pressure balance through density changes. White (2000) however, found that the SSH signal lags the SST signal by about $\pi/2$. He explained this predominantly in terms of an advective effect; however, the thermal signal may affect the surface wind, leading to a different phase relationship through ocean-atmosphere coupling.

Three mechanisms have been advocated to explain the signal in ocean colour. First, that they are simply due to north-south movements of water in a region of marked meridional gradient of CC. Second, that vertical adjustments within the water column have brought the deep phytoplankton up into the surface layer sensed by the satellite. However, simple calculations suggest this effect should be negligible. Third, and most

intriguingly, that the Rossby wave has affected the nutrient supply, leading to a genuine biological increase. [This is akin to phytoplankton growth in the core of an eddy, which was suggested by McGillicuddy et al., (1998) to be due to upwelling of nutrients.] To test these hypotheses, we review the CC observations in the light of those from other sensors.

a. Correspondence of individual events

In Fig. 1c we have highlighted a number of the westward-propagating local reductions in CC, and replicated these lines in Figs. 1b & 1d. The propagation speeds of these features (given by the slopes of the lines) vary from 1.2 to 4.2 km day⁻¹. Some of the features appear to undergo marked changes in speed or else generate secondary features of a different speed to the original. The different speeds are associated with different modes of Rossby waves, each with their own depth structure, and the changes in speed (and generation of new modes) occur where the bathymetry is varying most rapidly. Subrahmanyam et al. (2001) have shown that the first three modes of Rossby waves can be found in SSH data for the Indian Ocean, once a number of different latitudes is considered.

The majority of the highlighted features can be found as local increases in SST. Corresponding signals in SSH are not so readily apparent. Most of the *fast* features match a change in SSH so that the SST anomaly leads SSH by $\pi/2$, which leads CC by the same amount. In the east, the slower signals (with the more complex depth-structure) appear to have no analogue in SSH.

b. Correlations in phase and changes in amplitudes

As the closest agreement in the Hovmöller plots appears to be between SST and CC, we performed a lagged cross-correlation of the two. For the signals at 34°S, the strongest agreement was at zero lag, with a correlation of -0.45. Similar results are found for analyses of other sections at latitudes between 31° and 36°S; whereas further north the peak correlation (again at zero lag) is positive (see Fig. 2). As the SST always increases equatorwards (Fig. 3a), whereas the CC has a minimum at mid-latitudes, the observed change in sign of correlation coefficient (Fig. 2) may be thought consistent with where the SST and CC meridional gradients are of opposite or same sign. However the very broad minimum in the annual CC distribution does not produce significant gradients between 21°S and 29°S, whereas in that band there is sufficient CC signal to correlate positively with SST.

Clearly annual averages do not reveal the whole picture, and Fig. 1c illustrates the strength of seasonal modulation of the CC signal of Rossby waves at 34°S. The majority of the south Indian Ocean has a clear annual cycle in phytoplankton growth, with a peak in CC around September, and the sharp meridional gradient in CC in the south reaches its northernmost position in August (cf. Fig. 3b). However in the western half of the basin, between 30°S and 20°S, there is a secondary bloom in February and March (Longhurst, 2001) giving rise to strong short-lived CC gradients of opposing signs (see Fig. 3b). Advection cannot explain why the CC and SST anomalies appear in phase throughout the region north of 29°S (Fig. 2); thus other mechanisms may be playing an important rôle here.

The SST gradient at 34°S is $\sim 0.65^{\circ}\text{C} / \text{deg.}$ throughout the year, and the propagating signal has an amplitude of $\sim 0.6^{\circ}\text{C}$; for January to March, the CC gradient is about $-20\% / \text{deg.}$ and the propagating amplitude (averaged over 50°-105°E) is $\sim 20\%$. Thus the ratio of amplitude to gradient is the same for both CC and SST, giving support to the hypothesis that they are both acting as passive tracers in this region. Cipollini et al. (2000) compared the amplitudes and speeds of the various propagating features observed in the SSH and CC datasets. Neither showed significant eastward propagation. The strongest signature of Rossby waves in SSH was between 20° and 28°S, whereas that region had weak signals in the visible. The visible signal was most marked in the bands 11°-17°S and 31°-33°S, which is consistent with the strong meridional gradients in Fig. 3b. South of 20°S, the two datasets agreed on the dominant speed of signals; but further north, the ocean colour signal corresponded to a slower (higher order) mode than the SSH. Here, we note a similar mismatch at 34°S in the eastern side of the basin (Figs. 1c, d).

4. Conclusions

Rossby waves are known to be present in all ocean basins at most latitudes, but there are still many questions being raised concerning their propagation rates and the interchange of energy between the different modes. Although the mechanisms behind the ocean colour signal of Rossby waves are still being propounded and tested, it is clear that the combining of different observing techniques will greatly help address these questions.

Acknowledgements

The ATSR data were provided by the Rutherford Appleton Laboratory, SeaWiFS from NASA Goddard, and TOPEX/Poseidon data from AVISO. We thank all of these and the space agencies (ESA, NASA and CNES).

References

- Campbell, J. W., 1995: The lognormal distribution as a model for bio-optical variability in the sea. *Journal of Geophysical Research*, **100**, 13237-13254.
- Chelton, D. B. and M. G. Schlax, 1996: Global observations of oceanic Rossby waves. *Science*, **272**, 234-238.
- Cheney, R., L. Miller, R. Agreen, N. Doyle, J. Lillibridge, 1994: TOPEX/Poseidon: The 2cm solution. *Journal of Geophysical Research*, **99**, 24555-24563.
- Cipollini, P., D. Cromwell, P. G. Challenor, and S. Raffaglio, 2001: Rossby waves detected in global ocean colour data. *Geophysical Research Letters*, **28**, 323-326.
- Cipollini, P., D. Cromwell, M. S. Jones, G. D. Quartly, and P. G. Challenor, 1997: Concurrent altimeter and infrared observations of Rossby wave propagation near 34°N in the northeast Atlantic. *Geophysical Research Letters*, **24**, 889-892.
- Cipollini, P., P. G. Challenor, D. Cromwell, G. D. Quartly, and S. Raffaglio, 2000: Observation of Rossby waves in satellite derived chlorophyll-a data. *Proceedings of PORSEC '00*, Goa, India, Vol. II, 530-533.
- Gill, A. E., 1982: *Atmosphere-Ocean dynamics*, Academic Press, San Diego.
- Hill, K. L., I. S. Robinson, and P. Cipollini, 2000: Propagation characteristics of extratropical planetary waves observed in the ATSR global sea surface temperature record. *Journal of Geophysical Research*, **105**, 21927-21945.

- Hooker, S. B. and C. R. McClain., 2000: The calibration and validation of SeaWiFS data. *Progress in Oceanography*, **45**, 427-465.
- Longhurst, A., 2001: A major seasonal phytoplankton bloom in the Madagascar Basin. *Deep-Sea Research*, **I 48**, 2413-2422.
- McGillicuddy, D. J., A. R. Robinson, D. A. Siegel, H. W. Jannasch, R. Johnson, T. D. Dickey, J. McNeil, A. F. Michaels, and A. H. Knap, 1998: Influence of mesoscale eddies on new production in the Sargasso Sea. *Nature*, **394**, 263-266.
- Mutlow, C. T., A. M. Závody, I. J. Barton, and D. T. Llewellyn-Jones, 1994: Sea surface temperature measurements by the along-track scanning radiometer on the ERS-1 satellite: Early results. *Journal of Geophysical Research*, **99**, 22575-22588.
- Subrahmanyam, B., I. S. Robinson, J. R. Blundell, and P. G. Challenor, 2001: Indian Ocean Rossby waves observed in TOPEX/POSEIDON altimeter data and in model simulations. *International Journal of Remote Sensing*, **22**, 141-167.
- Tokmakian, R. T., and P. G. Challenor, 1993: Observations in the Canary Basin and the Azores Frontal region using Geosat data. *Journal of Geophysical Research*, **98**, 4761-4773.
- Uz, B. M., J. A. Yoder, and V. Osiachny, 2001: Pumping of nutrients to ocean surface waters by the action of propagating planetary waves. *Nature*, **409**, 597-600.
- White, W. B., 2000: Tropical coupled Rossby waves in the Pacific Ocean-Atmosphere system. *Journal of Physical Oceanography*, **30**, 1245-1264.

Figure captions

Figure 1: Hovmöller diagrams at 34°S in the Indian Ocean of various water properties. a) SST, b) SST anomalies, c) CC anomalies, d) SSH anomalies (see text for further details). e) Bathymetry for the section. [The solid lines are discussed in section 3a.]

Figure 2: Correlation of chlorophyll and temperature anomalies in the Indian Ocean as a function of latitude.

Figure 3: Mean meridional gradients between 50°E and 105°E. a) SST (in degs C per degree of latitude), b) Chlorophyll concentration (percentage change per degree of latitude).

Short Title

Rosby waves: Synergy in action




Figure 1: Hovmöller diagrams at 34°S in the Indian Ocean of various water properties. a) SST, b) SST anomalies, c) CC anomalies, d) SSH anomalies (see text for further details). e) Bathymetry for the section. [The solid lines are discussed in section 3a.]




Figure 2: Correlation of chlorophyll and temperature anomalies in the Indian Ocean as a function of latitude.




Figure 3: Mean meridional gradients between 50°E and 105°E. a) SST (in degs C per degree of latitude), b) Chlorophyll concentration (percentage change per degree of latitude).